

Structural plasticity and catalysis regulation of a thermosensor histidine kinase

Daniela Albanesi^{a,b,1}, Mariana Martín^{b,1}, Felipe Trajtenberg^c, María C. Mansilla^b, Ahmed Haouz^d, Pedro M. Alzari^a, Diego de Mendoza^b, and Alejandro Buschiazzi^{a,c,2}

^aInstitut Pasteur, Unité de Biochimie Structurale and ^dPlateforme de Cristallogénèse et Diffraction de Rayons X, URA 2185 Centre National de la Recherche Scientifique, Paris 75015, France; ^bInstituto de Biología Molecular y Celular de Rosario, Departamento de Microbiología, Facultad de Ciencias Bioquímicas y Farmacéuticas, Universidad Nacional de Rosario, Rosario 2000, Argentina; and ^cInstitut Pasteur de Montevideo, Unidad de Cristalografía de Proteínas, Montevideo 11400, Uruguay

Edited by E. Peter Greenberg, University of Washington School of Medicine, Seattle, WA, and approved August 3, 2009 (received for review June 15, 2009)

Temperature sensing is essential for the survival of living cells. A major challenge is to understand how a biological thermometer processes thermal information to optimize cellular functions. Using structural and biochemical approaches, we show that the thermosensitive histidine kinase, DesK, from *Bacillus subtilis* is cold-activated through specific interhelical rearrangements in its central four-helix bundle domain. As revealed by the crystal structures of DesK in different functional states, the plasticity of this helical domain influences the catalytic activities of the protein, either by modifying the mobility of the ATP-binding domains for autokinase activity or by modulating binding of the cognate response regulator to sustain the phosphotransferase and phosphatase activities. The structural and biochemical data suggest a model in which the transmembrane sensor domain of DesK promotes these structural changes through conformational signals transmitted by the membrane-connecting two-helical coiled-coil, ultimately controlling the alternation between output autokinase and phosphatase activities. The structural comparison of the different DesK variants indicates that incoming signals can take the form of helix rotations and asymmetric helical bends similar to those reported for other sensing systems, suggesting that a similar switching mechanism could be operational in a wide range of sensor histidine kinases.

coiled-coil | conformational rearrangement | crystallography | signal transduction

Cold sensors are ubiquitous integral membrane proteins found in all kingdoms of life. They are involved in many physiological roles, including membrane remodeling, chemotaxis, touch, and pain (1–3). The histidine kinase (HK), DesK, from *Bacillus subtilis* is the founding example of a membrane-bound thermosensor suited to remodel membrane fluidity when the ambient temperature drops below approximately 30 °C (Fig. S1). Several lines of evidence show that the cold thermal stimulus is detected by DesK, which together with the response regulator (RR), DesR, constitutes a canonical two-component system (TCS). In vivo experiments have demonstrated that DesK acts as a kinase at cold temperatures (4), ultimately activating the transcription of the gene *des* coding for the acyl lipid desaturase $\Delta 5$ -Des (1, 5). The increased fraction of unsaturated fatty acids in the membrane then restores fluidity and shuts off the kinase activity of DesK, terminating transcription.

Histidine kinases are multifunctional enzymes that share a conserved intracellular catalytic core linked to a high diversity of signal-sensing domains. Through still poorly understood mechanisms, HKs are able to catalyze autokinase, phosphotransferase, and protein phosphatase reactions in response to external stimuli, ultimately controlling the degree of phosphorylation of their cognate RR and hence the functional outcome of the signaling pathway. DesK is a class I HK (6) with an N-terminal sensor domain (≈ 150 residues) composed of four or five transmembrane (TM) segments connected to a C-terminal cytoplasmic domain (DesKC, ≈ 220 residues) (4) that belongs to the HisKA_3 subfamily (PFAM

00730) of HKs (7). To investigate how fluctuations in ambient temperature regulate the catalytic activities of DesK, we solved the crystal structure of its catalytic core in different functional states and determined the functional properties of the full-length sensor in pure lipids vesicles. The results highlight the remarkable plasticity of the central four-helix bundle domain as the protein proceeds along the catalytic cycle, and suggest a signal-dependent regulation model that may be operational in a wide range of HKs.

Results

To uncover the structural features of DesK associated with the different functional states of the protein, we carried out structural studies of DesKC before and after autophosphorylation. We also characterized two point mutants of the catalytic domain, in which the phosphorylatable histidine residue (H188) was substituted either by valine (DesKC_{H188V}), a mutant known to retain the phosphatase activity of the wild-type protein (4), or by glutamic acid (DesKC_{H188E}), sought to mimic the phosphorylated state of the enzyme. The 3D structures of the different DesKC variants were determined in six different crystalline environments (Table 1) using multiwavelength anomalous diffraction (MAD) and molecular replacement methods (SI Text and Table S1).

The catalytic core of DesK (Fig. 1A–C) shows the characteristic homodimeric structure observed in other HKs (8). Each monomer consists of an N-terminal antiparallel 2-helix hairpin (helices $\alpha 1$ and $\alpha 2$) that includes the phosphorylatable H188, connected by a short linker region (residues 243–245) to a C-terminal ATP-binding domain (ABD). The helical hairpins of two monomers interact with each other to form a central four-helix bundle (4-HB) domain, known as the DHp (for Dimerization and Histidine phosphotransfer) domain. In each monomer, the N-terminal part of helix $\alpha 1$ extends beyond the 4-HB, connecting the catalytic core with the TM sensor domain. The structure of the ABD, composed by a five-stranded β -sheet opposed by a layer of three α -helices, is similar to that of other members of the GHKL superfamily (9) and remains essentially unchanged in the different DesKC structures (Fig. 1D), except for a high mobility of the ATP lid.

Three Distinct Conformational States of DesKC Variants. The structures of DesKC reveal three distinct conformational states of the

Author contributions: A.H., P.M.A., D.d.M., and A.B. designed research; D.A., M.M., F.T., A.H., P.M.A., D.d.M., and A.B. performed research; D.A., M.M., F.T., M.C.M., and A.H. contributed new reagents/analytic tools; D.A., M.M., F.T., M.C.M., P.M.A., D.d.M., and A.B. analyzed data; and P.M.A., D.d.M., and A.B. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: The atomic coordinates and structure factors have been deposited in the Protein Data Bank, www.pdb.org (PDB ID codes 3EHF, 3EHG, 3EHJ, 3GIE, 3GIF, and 3GIG).

¹D.A. and M.M. contributed equally to this work.

²To whom correspondence should be addressed. E-mail: alebus@pasteur.edu.uy.

This article contains supporting information online at www.pnas.org/cgi/content/full/0906699106/DCSupplemental.

a (K^+/P^+) phenotype. Upon cold signal reception, the ensuing structural reorganization would release the ABDs for histidine phosphorylation, as seen in DesKC $_{\Delta 174}$ (Fig. 6, *Center*). Compelling mutagenesis evidence supporting the importance of weakening the DHP-ABD association for autokinase activity has been previously obtained for TM0853 (13). In this functional state, two distinct factors contribute to down-regulate DesK phosphatase activity. First, the modified DHP interaction surface precludes DesK/DesR binding, as suggested by the lack of complex formation of the unphosphorylated proteins (Fig. S6C). Second, the dynamic effect of the released ABDs, covalently attached to the DHP domain, could also contribute to decrease the resultant DesR-binding affinity due to a higher effective concentration. These mechanisms (surface modulation, ABD mobility) are not mutually exclusive and are amenable to fine-tuning in different HKs. Phosphorylation of DesKC induces an as yet different, asymmetric conformation with a pronounced bending of helix $\alpha 1$ (Fig. 6, *Right*), capable of interacting with DesR (Fig. S6B). Crystallization of DesKC $_{H188E}$ in two distinct conformations, respectively similar to DesKC $_{\Delta 174}$ (autokinase) and DesKC-P (phosphotransferase), suggest a low inter-conversion energy barrier between these functional states. As a consequence, the corresponding structural transition could be induced, in the absence of external stimuli, by phosphate attachment to or transfer from H188 (as proposed in Fig. 6, *Lower*).

The proposed model of catalysis regulation in DesK, governed by interhelical rearrangements in the DHP coiled-coil, may be a common trait for a large fraction of HKs. Alpha helices are indeed common information transducer elements (28) and different types of helical movements performing mechanical work have been proposed to convey input/output signals (29–31). Furthermore, incoming signals in the form of helix rotations are congruent with mechanistic models derived from the study of upstream elements in different signal transducing systems. Thus, the parallel coiled-coil HAMP domain, present in the membrane-connecting region of several HKs (32), was shown to relay the input signal through helical rotations (25) compatible with the structural changes seen in DesK. Also, a combined helical rotation and tilting was involved in signal transduction by the bacterial sensory rhodopsin phototactic receptor (33) and analogous rotational movements were recently proposed as a general mechanism of signal transduction through α -helical coiled-coil linkers (34).

Materials and Methods

Protein Production and Crystallization. All recombinant proteins were produced and purified as described in *SI Text*. For DesKC phosphorylation, the protein (10 mg/mL) was preincubated in 50 mM Tris-HCl, pH 8.5, with 5 mM ATP and 10 mM MgCl $_2$ for 1 h at room temperature. All proteins were preincubated with 5 mM AMP-PCP or ADP (as indicated for each structure in Table S1) before crystallization. Initial crystallogenic screenings were performed as described in *SI Text*. After optimization, crystals were grown at 18 °C by mixing 2 mL protein solution (10 mg/mL) and 2 mL reservoir solution in hanging drops. Reservoir solutions contained: 10% PEG-3350, 0.1 M Bicine, pH 9, 0.1 M CaCl $_2$, and 30% glycerol (V188a, V188b); 18% PEG-8000, 0.1 M Mes, pH 6.5, and 0.2 M calcium acetate (DesKC $_{\Delta 174}$); 10% PEG 3000, 0.1 M CHES, pH 9.5, and 10 mM MgCl $_2$ (DesKC-P, 7 mg/mL); 14% MPD, 0.1 M HEPES, pH 6, and 10 mM MgCl $_2$ (E188b); and 14% PEG-3350, 0.1 M Tris, pH 7, 5 mM KCl, and 50 mM CaCl $_2$ (E188a).

Crystallographic Studies. Single crystals were cryoprotected in mother liquor containing 25% glycerol (or 22% MPD) and flash frozen in liquid N $_2$ before X-ray data collection; all diffraction data sets were processed according to standard procedures (full details in *SI Text* and Table S1). The structures were solved by multiwavelength anomalous diffraction (MAD), using SeMet-labeled proteins, except for DesKC-P and DesKC $_{H188E}$, which were solved by molecular replacement methods. Complete procedures of data processing, selenium substructure determination and protein phasing/refinement are detailed in *SI Text*; final refinement parameters are given in Table S1.

Size-Exclusion Chromatography. Proteins (1–1.5 mg/mL) were preincubated with 10 mM MgCl $_2$ and 5 mM ADP or ATP for 30 min and then injected in a Superdex75 10/300 column (GE Healthcare), equilibrated in 50 mM HEPES, pH 7.5, 0.3 M NaCl, and 10 mM MgCl $_2$ and run at 0.5 mL/min.

Proteoliposomes Obtainment, Purification, and Activity Characterization. The synthesis and purification of DesK proteoliposomes and the enzyme activity assays were performed following the published method (23), with the modifications described in *SI Text*.

ACKNOWLEDGMENTS. We thank the staff of the Unidad de Bioquímica Analítica (Inst Pasteur Montevideo) for mass spectrometry analyses and of beamlines ID23-1, ID14-4 and ID-29 (European Synchrotron Radiation Facility) for diffraction data collection. This work was supported by grants from Agencia Nacional de Investigación e Innovación, Uruguay; Agence Nationale de la Recherche, France; Institut Pasteur (France); Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina; Agencia de Promoción Científica y Tecnológica, Argentina; and ECOS France-Argentina Action Grant A05B02. D.A. is a Marie Curie Incoming International Fellow (European Union), F.T. is a fellow from Agencia Nacional de Investigación e Innovación, and D.d.M. is an International Research Scholar from the Howard Hughes Medical Institute.

- Aguilar PS, Hernandez-Arriaga AM, Cybulski LE, Erazo AC, de Mendoza D (2001) Molecular basis of thermosensing: A two-component signal transduction thermometer in *Bacillus subtilis*. *EMBO J* 20:1681–1691.
- Dhaka A, et al. (2007) TRPM8 is required for cold sensation in mice. *Neuron* 54:371–378.
- Salman H, Libchaber A (2007) A concentration-dependent switch in the bacterial response to temperature. *Nat Cell Biol* 9:1098–1100.
- Albanesi D, Mansilla MC, de Mendoza D (2004) The membrane fluidity sensor DesK of *Bacillus subtilis* controls the signal decay of its cognate response regulator. *J Bacteriol* 186:2655–2663.
- Cybulski LE, del Solar G, Craig PO, Espinosa M, de Mendoza D (2004) *Bacillus subtilis* DesR functions as a phosphorylation-activated switch to control membrane lipid fluidity. *J Biol Chem* 279:39340–39347.
- Bilwes AM, Alex LA, Crane BR, Simon MI (1999) Structure of CheA, a signal-transducing histidine kinase. *Cell* 96:131–141.
- Finn RD, et al. (2008) The Pfam protein families database. *Nucleic Acids Res* 36:D281–288.
- Varughese KI, Madhusudan, Zhou XZ, Whiteley JM, Hoch JA (1998) Formation of a novel four-helix bundle and molecular recognition sites by dimerization of a response regulator phosphotransferase. *Mol Cell* 2:485–493.
- Dutta R, Inouye M (2000) GHKL, an emergent ATPase/kinase superfamily. *Trends Biochem Sci* 25:24–28.
- Walshaw J, Woolfson DN (2001) Open-and-shut cases in coiled-coil assembly: Alpha-sheets and alpha-cylinders. *Protein Sci* 10:668–673.
- Crick F (1953) The packing of alpha-helices: Simple coiled-coils. *Acta Crystallogr* 6:689–697.
- Lupas AN, Gruber M (2005) The structure of alpha-helical coiled coils. *Adv Protein Chem* 70:37–78.
- Marina A, Waldburger CD, Hendrickson WA (2005) Structure of the entire cytoplasmic portion of a sensor histidine-kinase protein. *EMBO J* 24:4247–4259.
- Bick MJ, et al. (2009) How to switch off a histidine kinase: Crystal structure of *Geobacillus stearothermophilus* KinB with the inhibitor Sda. *J Mol Biol* 386:163–177.
- Cai SJ, Inouye M (2003) Spontaneous subunit exchange and biochemical evidence for trans-autophosphorylation in a dimer of *Escherichia coli* histidine kinase (EnvZ). *J Mol Biol* 329:495–503.
- Park H, Saha SK, Inouye M (1998) Two-domain reconstitution of a functional protein histidine kinase. *Proc Natl Acad Sci USA* 95:6728–6732.
- Tanaka T, et al. (1998) NMR structure of the histidine kinase domain of the *E. coli* osmosensor EnvZ. *Nature* 396:88–92.
- Neiditch MB, et al. (2006) Ligand-induced asymmetry in histidine sensor kinase complex regulates quorum sensing. *Cell* 126:1095–1108.
- Pollard AM, Bilwes AM, Crane BR (2009) The structure of a soluble chemoreceptor suggests a mechanism for propagating conformational signals. *Biochemistry* 48:1936–1944.
- Hsing W, Russo FD, Bernd KK, Silhavy TJ (1998) Mutations that alter the kinase and phosphatase activities of the two-component sensor EnvZ. *J Bacteriol* 180:4538–4546.
- Zhu Y, Qin L, Yoshida T, Inouye M (2000) Phosphatase activity of histidine kinase EnvZ without kinase catalytic domain. *Proc Natl Acad Sci USA* 97:7808–7813.
- Cybulski LE, et al. (2002) Mechanism of membrane fluidity optimization: Isothermal control of the *Bacillus subtilis* acyl-lipid desaturase. *Mol Microbiol* 45:1379–1388.
- Martin M, Albanesi D, Alzari PM, de Mendoza D (2009) Functional in vitro assembly of the integral membrane bacterial thermosensor DesK. *Protein Expr Purif* 66:39–45.
- Cronan JE, Jr, Gelmann EP (1975) Physical properties of membrane lipids: Biological relevance and regulation. *Bacteriol Rev* 39:232–256.
- Hulko M, et al. (2006) The HAMP domain structure implies helix rotation in transmembrane signaling. *Cell* 126:929–940.
- Deng Y, et al. (2006) Antiparallel four-stranded coiled coil specified by a 3–3–1 hydrophobic heptad repeat. *Structure* 14:247–255.
- Yadav MK, et al. (2006) Coiled coils at the edge of configurational heterogeneity: Structural analyses of parallel and antiparallel homotetrameric coiled coils reveal configurational sensitivity to a single solvent-exposed amino acid substitution. *Biochemistry* 45:4463–4473.
- Strickland D, Moffat K, Sosnick TR (2008) Light-activated DNA binding in a designed allosteric protein. *Proc Natl Acad Sci USA* 105:10709–10714.
- Gao R, Lynn DG (2007) Integration of rotation and piston motions in coiled-coil signal transduction. *J Bacteriol* 189:6048–6056.
- Kwon O, Georgellis D, Lin EC (2003) Rotational on-off switching of a hybrid membrane sensor kinase Tar-ArcB in *Escherichia coli*. *J Biol Chem* 278:13192–13195.
- Ottemann KM, Xiao W, Shin YK, Koshland DE, Jr (1999) A piston model for transmembrane signaling of the aspartate receptor. *Science* 285:1751–1754.
- Szurmant H, White RA, Hoch JA (2007) Sensor complexes regulating two-component signal transduction. *Curr Opin Struct Biol* 17:706–715.
- Moukhametzanov R, et al. (2006) Development of the signal in sensory rhodopsin and its transfer to the cognate transducer. *Nature* 440:115–119.
- Moglich A, Ayers RA, Moffat K (2009) Design and signaling mechanism of light-regulated histidine kinases. *J Mol Biol* 385:1433–1444.